arXiv:1302.1441v1 [math.AC] 6 Feb 2013

ADDENDUM TO UNIQUENESS OF CERTAIN POLYNOMIALS CONSTANT ON A LINE

JIŘÍ LEBL

Abstract. The computer calculations in [2] to classify sharp polynomials with nonegative coefficients constant on the line $x + y = 1$ have been extended to degree 19, where a surprisingly large number of 13 sharp polynomials was found.

1. Introduction

Let $\mathcal{H}(2, d)$ be the space of polynomials $p(x, y)$ of two variables with nonnegative coefficients such that $p(x, y) = 1$ whenever $x + y = 1$. It is known [1] that the degree $d \leq 2N - 3$, where N is the number of positive coefficients of p. Furthermore, it is known that for each odd degree d, there exists a group invariant polynomial with precisely $d = 2N - 3$. We call such polynomials the *sharp polynomials*. See [1, 2] for more information, background, and motivation.

In [2] using computer code, we have have classified the sharp polynomials in $\mathcal{H}(2, d)$ up to degree 17. Due to the increase in the speed of computers over the last several years and improvements to the computer code to be described below, it was possible to run the computer code for degree 19. The computation with all the improvements takes approximately 5 days on 4 relatively recent CPU cores.

The computer has found 13 polynomials polynomials in degree 19 up to symmetry of the variables, several of which were unexpected. Two of the polynomials are symmetric. The number of sharp polynomials in degree 19 is stunning as there have been only 16 sharp polynomials in all odd degrees up to degree 17.

We have therefore computed a new term for the sequence A143107 on OEIS [4]. That is a sequence whose Nth term is the number of sharp polynomials of degree $2N - 3$, or in other words, the number of polynomials in $\mathcal{H}(2, 2N - 3)$ with N nonzero coefficients. In this sequence symmetry is not taken into account and therefore there are 24 polynomials altogether for degree 19. The sequence is now known to be:

$$0, 1, 1, 2, 4, 2, 4, 8, 4, 2, 24. \tag{1}$$

At the time of this writing, the code is running for degree 21, and it is expected to finish in approximately a year. We expect that only the group invariant polynomial is found in degree 21, and so we expect the next term in the sequence to be 2.

Date: February 7, 2013.

The first author was in part supported by NSF grant DMS 0900885.

2. The polynomials

Let us list the polynomials that were found. First we have the group invariant polynomial.

$$\begin{aligned} &x^{19} + 19\,y\,x^{17} + 152\,y^2\,x^{15} + 665\,y^3\,x^{13} + 1729\,y^4\,x^{11} + 2717\,y^5\,x^9 + 2508\,y^6\,x^7 + \\ &\qquad 1254\,y^7\,x^5 + 285\,y^8\,x^3 + 19\,y^9\,x + y^{19} \end{aligned} \tag{2}$$

Then we have several polynomials with integer coefficients.

$$\begin{aligned} &x^{19} + 19\,y\,x^{17} + 152\,y^2\,x^{15} + 665\,y^3\,x^{13} + 1729\,y^4\,x^{11} + 2090\,y^5\,x^9 + 627\,y^9\,x^5 + \\ &\qquad 627\,y^5\,x^5 + 285\,y^8\,x^3 + 19\,y^9\,x + y^{19} \end{aligned} \tag{3}$$

$$\begin{aligned} &x^{19} + 285\,y^3\,x^{13} + 1425\,y^4\,x^{11} + 19\,y^9\,x^9 + 2679\,y^5\,x^9 + 19\,y\,x^9 + 2508\,y^6\,x^7 + \\ &\qquad 1254\,y^7\,x^5 + 285\,y^8\,x^3 + 19\,y^9\,x + y^{19} \end{aligned} \tag{4}$$

$$\begin{aligned} &x^{19} + 285\,y^3\,x^{13} + 1425\,y^4\,x^{11} + 19\,y^9\,x^9 + 2052\,y^5\,x^9 + 19\,y\,x^9 + 627\,y^9\,x^5 + \\ &\qquad 627\,y^5\,x^5 + 285\,y^8\,x^3 + 19\,y^9\,x + y^{19} \end{aligned} \tag{5}$$

Next, three polynomials with rational coefficients with denominator 25.

$$\begin{aligned} &x^{19} + 19\,y\,x^{17} + 152\,y^2\,x^{15} + \frac{15371\,y^3\,x^{13}}{25} + \frac{6137\,y^4\,x^{11}}{5} + \frac{4807\,y^5\,x^9}{5} + \\ &\qquad \frac{1254\,y^{13}\,x^3}{25} + \frac{4617\,y^8\,x^3}{25} + \frac{1254\,y^3\,x^3}{25} + 19\,y^9\,x + y^{19} \end{aligned} \tag{6}$$

$$\begin{aligned} &x^{19} + \frac{5871\,y^3\,x^{13}}{25} + \frac{4617\,y^4\,x^{11}}{5} + 19\,y^9\,x^9 + \frac{4617\,y^5\,x^9}{5} + 19\,y\,x^9 + \\ &\qquad \frac{1254\,y^{13}\,x^3}{25} + \frac{4617\,y^8\,x^3}{25} + \frac{1254\,y^3\,x^3}{25} + 19\,y^9\,x + y^{19} \end{aligned} \tag{7}$$

$$\begin{aligned} &x^{19} + \frac{5871\,y^3\,x^{13}}{25} + \frac{4617\,y^4\,x^{10}}{5} + 19\,y^9\,x^9 + \frac{4617\,y^6\,x^9}{5} + 19\,y\,x^9 + \\ &\qquad \frac{1254\,y^{13}\,x^3}{25} + \frac{4617\,y^8\,x^3}{25} + \frac{1254\,y^3\,x^3}{25} + 19\,y^9\,x + y^{19} \end{aligned} \tag{8}$$

Then we have two polynomials with denominator 56.

$$\begin{aligned} &x^{19} + \frac{855\,y\,x^{17}}{56} + \frac{646\,y^2\,x^{15}}{7} + \frac{1938\,y^3\,x^{13}}{7} + \frac{2907\,y^4\,x^{11}}{7} + \\ &\qquad \frac{3553\,y^5\,x^9}{14} + \frac{323\,y^8\,x^3}{7} + \frac{209\,y^{17}\,x}{56} + \frac{323\,y^9\,x}{28} + \frac{209\,y\,x}{56} + y^{19} \end{aligned} \tag{9}$$

$$\begin{aligned} &x^{19} + \frac{855\,y\,x^{17}}{56} + \frac{323\,y^2\,x^{15}}{7} + \frac{323\,y^8\,x^9}{7} + \frac{323\,y^5\,x^9}{2} + \\ &\qquad \frac{323\,y^2\,x^9}{7} + \frac{323\,y^8\,x^3}{7} + \frac{209\,y^{17}\,x}{56} + \frac{323\,y^9\,x}{28} + \frac{209\,y\,x}{56} + y^{19} \end{aligned} \tag{10}$$

Finally, rather surprisingly, there are 4 very similar polynomials with denominator 110, two of which are symmetric in x and y.

$$\begin{multline} x^{19}+\frac{19\,y\,x^{17}}{2}+\frac{323\,y^2\,x^{15}}{11}+\frac{323\,y^3\,x^{13}}{11}+\frac{323\,y\,x^6}{55}+ \\ \frac{323\,y^{13}\,x^4}{11}+\frac{323\,y^{14}\,x^2}{11}+\frac{19\,y^{17}\,x}{2}+\frac{323\,y^6\,x}{55}+\frac{399\,y\,x}{110}+y^{19} \tag{11} \end{multline}$$

$$\begin{multline} x^{19}+\frac{19\,y\,x^{17}}{2}+\frac{323\,y^2\,x^{15}}{11}+\frac{323\,y^3\,x^{13}}{11}+\frac{323\,y\,x^6}{55}+ \\ \frac{323\,y^{13}\,x^3}{11}+\frac{323\,y^{15}\,x^2}{11}+\frac{19\,y^{17}\,x}{2}+\frac{323\,y^6\,x}{55}+\frac{399\,y\,x}{110}+y^{19} \tag{12} \end{multline}$$

$$\begin{multline} x^{19}+\frac{19\,y\,x^{17}}{2}+\frac{323\,y^2\,x^{14}}{11}+\frac{323\,y^4\,x^{13}}{11}+\frac{323\,y\,x^6}{55}+ \\ \frac{323\,y^{13}\,x^4}{11}+\frac{323\,y^{14}\,x^2}{11}+\frac{19\,y^{17}\,x}{2}+\frac{323\,y^6\,x}{55}+\frac{399\,y\,x}{110}+y^{19} \tag{13} \end{multline}$$

$$\begin{multline} x^{19}+\frac{19\,y\,x^{17}}{2}+\frac{323\,y^2\,x^{14}}{11}+\frac{323\,y^4\,x^{13}}{11}+\frac{323\,y\,x^6}{55}+ \\ \frac{323\,y^{13}\,x^3}{11}+\frac{323\,y^{15}\,x^2}{11}+\frac{19\,y^{17}\,x}{2}+\frac{323\,y^6\,x}{55}+\frac{399\,y\,x}{110}+y^{19} \tag{14} \end{multline}$$

3. The computer code

The code used is the C code described in detail in [2]. The new revision of the code that was used in this computation has been posted at [3]. The basic idea is to consider $p(x, 1-x) - 1 = 0$, which provides a linear equation for the coefficients. We write this equation as a matrix A that takes coefficients of degree $d-1$ or less to the degree d coefficients. We know that the degree d coefficients have the form $x^d + y^d$. We iterate over the list of possible monomials of degree $d-1$ or less, taking the corresponding columns of the matrix, we look for nonzero solutions. The idea of the algorithm is to find those submatrices that are not of full rank (have a nontrivial solution). Then we check if this solution has positive coefficients. There are several heuristics that are applied that can avoid doing row reduction at all. Already in [2], to improve speed of the row reduction, we first used mod p arithmetic for reduction for a small prime, as most of the submatrices considered are full rank. This technique reduces the need to do row reduction in full integer arithmetic in vast majority of the cases. We have used $p = 19$ in degree 17 or less, and in this calculation, we used $p = 23$. The prime must not divide the degree, as most entries in the matrix are divisible by d.

The major improvements that were done in this revision the following

(i) Before mode p, the row reduction is first done mod 2. Mod 2 is much faster than mod p. The code actually does column reduction in mod 2 and considers columns as unsigned integers. The reduction removes one internal loop as adding columns together is simply an XOR operation. Unfortunately mod 2 arithmetic only eliminates 90% of the full rank cases, but it is approximately 4 times as fast on these 90%. If the test fails on mod 2, we move to mod p as before.

(ii) Just as the Mathematica code from [2], the C code now ignores polynomials that are "right side heavy." That is, polynomials where there are more terms with higher power of y than of x. This optimization reduces the run time by approximately one third.

(iii) Many other more minor optimizations were done, whose individual impact was harder to measure.

Overall, together with improvements in speed of computers since 2008, the current code runs approximately 25 times faster than it did in 2008.

References

[1] John P. D'Angelo, Šimon Kos, and Emily Riehl, *A sharp bound for the degree of proper monomial mappings between balls*, J. Geom. Anal. **13** (2003), no. 4, 581–593. MR2005154

[2] Jiří Lebl and Daniel Lichtblau, *Uniqueness of certain polynomials constant on a hyperplane*, Linear Algebra Appl. **433** (2010), no. 4, 824–837. arXiv:0808.0284. MR2654111

[3] Jiří Lebl, *Computer code used.* `http://www.jirka.org/rref-v2.zip`.

[4] OEIS Foundation Inc., *The on-line encyclopedia of integer sequences.* `http://oeis.org/A143107`.

Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA
E-mail address: `lebl@math.wisc.edu`